



On the cover: A jacketed Franz diffusion cell remains one of the most popular cells that scientists use when investigating transdermal drug permeation. The introduction of a reliable in vitro system has enabled a significant reduction in the number of in vivo experiments, lowering the cost and facilitating the identification of potential candidates for percutaneous drug delivery.

Courtesy of Mikolaj Milewski, a graduate student working with Audra Stinchcomb at the University of Kentucky College of Pharmacy, Lexington, Kentucky, USA. E-mail: mmile2@email.uky.edu

www.transdermalmag.com May 2010 Volume 2 Number 3

IN VITRO TESTING OF TRANSDERMALS

Human Skin Equivalent (HSEs) as Alternatives for Transdermal Permeation, Phototoxicity, and Cytotoxicity Studies 5

*Vishwas Rai, Jenna Terebetski, and B. Michniak-Kohn
Rutgers - The State University of New Jersey.*

API Heterogeneity in a Multilayered Transdermal Patch 9

Oksana Klueva, ChemImage

SKIN PERMEATION: CHEMICAL ENHANCEMENTS

Chemical Enhancements of Transdermal Drug Delivery Systems: Advantages and Challenges—Part 2 15

*Danyi Quan, Xel Pharmaceuticals, Inc, and
Xiaoying Hui and Howard I. Maibach, Department of Dermatology,
University of California San Francisco*

DRUG DELIVERY: FOLLICULAR ROUTE

The Follicular Route for Drug Delivery 21

Philip W. Wertz, University of Iowa

DEPARTMENTS

Industry news 3	Calendar Cover 3
Research news 4	Supplier news Cover 3
Issue Focus 13	People news Cover 3